arXiv:1610.03923v1 [math.DG] 13 Oct 2016

LIOUVILLE PROPERTY FOR f-HARMONIC FUNCTIONS WITH POLYNOMIAL GROWTH

JIA-YONG WU

Abstract. We prove a Liouville property for any f-harmonic function with polynomial growth on a complete noncompact smooth metric measure space $(M, g, e^{-f}dv)$ whose Bakry-Émery Ricci curvature satisfies a quadratic decay lower bound and whose any diameter of geodesic sphere has sublinear growth.

1. Introduction

In this paper, we will study a Liouville property for any f-harmonic function with polynomial growth on complete smooth metric measure spaces with lower quadratic Bakry-Émery Ricci curvature decay.

Recall that an n-dimensional complete smooth metric measure space, customarily denoted by $(M, g, e^{-f}dv_g)$, is an n-dimensional Riemannian manifold (M, g) together with a weighted volume element $e^{-f}dv_g$ for some $f \in C^\infty(M)$, and the volume element dv_g induced by the Riemannian metric g. The f-Laplacian Δ_f on smooth metric measure space $(M, g, e^{-f}dv)$, self-adjoint with respect to $e^{-f}dv_g$, is defined by

$$\Delta_f = \Delta - \nabla f \cdot \nabla.$$

On smooth metric measure space $(M, g, e^{-f}dv_g)$, a smooth function u is called weighted harmonic (or f-harmonic) if $\Delta_f u = 0$, and f-subharmonic if $\Delta_f u \geq 0$. It is easy to see that the absolute value of an f-harmonic function is a nonnegative f-subharmonic function. If f is constant, all notions mentioned above reduce to the classical Riemannian case.

A natural generalization of Ricci curvature associated to smooth metric measure space $(M, g, e^{-f}dv_g)$ is called m-Bakry-Émery Ricci curvature, which is defined by

$$\mathrm{Ric}_f^m = \mathrm{Ric} + \nabla^2 f - \frac{1}{m} df \otimes df$$

for some number $m > 0$. Bakry and Emery [1] extensively studied this curvature tensor and its relationship to diffusion processes. When m is

Date: July 8, 2016.

2000 *Mathematics Subject Classification.* Primary 53C21; Secondary 58J05.

Key words and phrases. Bakry-Émery Ricci curvature; smooth metric measure space; Liouville theorem.

Dedicated to my daughter Yuxuan for her third birthday.

finite, the Bochner formula for the m-Bakry-Émery Ricci curvature can be read as

$$\frac{1}{2}\Delta_f|\nabla u|^2 \geq \frac{(\Delta_f u)^2}{m+n} + \langle \nabla \Delta_f u, \nabla u\rangle + \mathrm{Ric}_f^m(\nabla u, \nabla u),$$

which could be viewed as the Bochner formula for the Ricci curvature of an $(n+m)$-dimensional manifold. Hence many classical results for manifolds with Ricci curvature bounded below can be extended to smooth metric measure spaces with m-Bakry-Émery Ricci curvature bounded below, see for example [2, 12, 17, 19, 24] for details. When $m=\infty$, one denotes

$$\mathrm{Ric}_f = \mathrm{Ric}_f^\infty = \mathrm{Ric} + \nabla^2 f.$$

In particular, if

$$\mathrm{Ric}_f = \lambda\, g$$

for some $\lambda \in \mathbb{R}$, then $(M, g, e^{-f}dv_g)$ is a gradient Ricci soliton, which arise from the singularity analysis of the Ricci flow [7]. When Ric_f is bounded below, some analytical and geometric properties on smooth metric measure spaces were also successfully explored as long as the weight function f is assumed to be some condition. We refer to the work of Lott [14], Wei and Wylie [18], and [20, 22, 23, 24] for further details.

Recently, many Liouville theorems for f-harmonic functions on smooth metric measure spaces have been studied. For example, Wei and Wylie [18] proved that any positive f-harmonic function with some growth condition with $\mathrm{Ric}_f \geq K > 0$ must be constant. Brighton [3] generalized Yau's result [26] and derived a gradient estimate for positive f-harmonic functions. He also obtained a Liouville theorem for positive bounded f-harmonic functions. Inspired by Brighton's argument, Munteanu and Wang [15] applied the De Giorgi-Nash-Moser theory to get a sharp gradient estimate and a Liouville-type result for positive f-harmonic functions on smooth metric measure spaces. X.-D. Li [13] applied probabilistic arguments to gave an alternative proof of Brighton's gradient estimate and Liouville theorem for positive f-harmonic functions. In [20], the author got a local sharp elliptic gradient estimate for positive solutions to the f-heat equation, using which he proved parabolic Liouville theorems for f-heat equation with certain growth restriction on smooth metric measure spaces when $\mathrm{Ric}_f \geq 0$.

Motivated by the classical works of [10, 11, 25], many L^p_f-Liouville ($0 < p < \infty$) theorems on smooth metric measure spaces were extensively studied. Here L^p_f-Liouville theorem means that every L^p_f-integrable f-harmonic function on smooth metric measure spaces is constant. Recall that u is called L^p_f-integrable, i.e. $u \in L^p_f$, if its L^p_f-norm, defined as

$$\|u\|_p := \left(\int_M |u|^p e^{-f} dv\right)^{1/p},$$

is finite. In [12] X.-D. Li proved an L^1_f-Liouville theorem for f-subharmonic functions on smooth metric measure spaces with $\mathrm{Ric}_f^m \geq -c(1+r(x)^2)$,

$0 < m < \infty$. Pigola, Rimoldi, and Setti [16] proved a sharp L^p_f-Liouville theorem with $p > 1$ for f-subharmonic functions on smooth metric measure spaces without any curvature assumption. When f is bounded, the author [20] derived some L^p_f-Liouville theorems with $0 < p \leq 1$ for f-subharmonic functions on smooth metric measure spaces with some Bakry-Emery curvature assumptions. When f is not bounded, P. Wu and the author [22, 23, 24] systematically studied various L^p_f-Liouville theorems with $0 < p \leq 1$ on smooth metric measure spaces under different Bakry-Emery curvature conditions. In particular, we proved a sharp L^1_f-Liouville theorem when $\mathrm{Ric}_f \geq 0$ by using f-heat kernel Gaussian estimates on smooth metric measure spaces (see also [5]).

In this paper, we prove a new Liouville theorem for any f-harmonic function with polynomial growth when Bakry-Émery Ricci curvature satisfies a quadratic decay lower bound and the diameter of geodesic sphere is sublinear. The proof uses a local f-Neumann Poincaré inequality, and an iterated argument combined with a cut off function technique, similar to the arguments of Carron in [4] (see also [22, 23]).

Theorem 1.1. *Let $(M, g, e^{-f}dv)$ be an n-dimensional complete noncompact smooth metric measure space. For a base point $o \in M$, assume that*

$$\mathrm{Ric}_f \geq -c\, r^{-2}(x)$$

for some constant $c > 0$, where $r(x) := d(o, x)$ is the distance function, and

$$\lim_{R\to\infty} \sup_{x\in B(o,R)} |f(x)| < +\infty.$$

Also if the diameter of geodesic sphere has a sublinear growth:

$$\mathrm{diam}\, \partial B(o, R) := \sup_{x,y\in\partial B(o,R)} d(x, y) = o(R), \quad R \to \infty,$$

then any f-harmonic function with polynomial growth is constant.

Remark 1.2. In the proof of Theorem 1.1, the finiteness of f at infinity makes the iterated procedure to work efficiently. It is interesting to ask whether this condition can be removed.

Remark 1.3. For smooth Riemannian manifolds whose sectional curvature satisfies a quadratic decay lower bound and whose geodesic spheres have sublinear growth, A. Kasue [8, 9] proved that any harmonic function with polynomial growth is constant (see also G. Carron [4] for further generalizations). Theorem 1.1 generalizes Kasue's result to the setting of smooth metric measure spaces.

This rest of the paper is organized as follows. In Section 2, we recall some known results, such as f-volume comparison theorems, local Poincaré inequalities for Bakry-Émery Ricci curvature bounded below. In Section 3, we adapt an iterated argument to prove Theorem 1.1.

Acknowledgement. This work was partially supported by the NSFC (11271132) and the Innovation Program of Shanghai Municipal Education Commission (13YZ087).

2. Preliminaries

Let $(M, g, e^{-f}dv)$ be an n-dimensional complete noncompact smooth metric measure space. For any point $x \in M$ and $R > 0$, we denote

$$A(R) := A(x, R) = \sup_{y \in B_x(3R)} |f(y)|.$$

We often write A for short. In [18] Wei and Wylie proved a relative f-volume comparison theorem.

Lemma 2.1. *Let $(M, g, e^{-f}dv)$ be an n-dimensional complete noncompact smooth metric measure space. If*

$$\mathrm{Ric}_f \geq -(n-1)\,K$$

for some constant $K \geq 0$, then

$$\frac{V_f(B_x(R_1, R_2))}{V_f(B_x(r_1, r_2))} \leq \frac{V_K^{n+4A}(B_x(R_1, R_2))}{V_K^{n+4A}(B_x(r_1, r_2))} \tag{2.1}$$

for any $0 < r_1 < r_2,\ 0 < R_1 < R_2,\ r_1 \leq R_1,\ r_2 \leq R_2$, where $B_x(R_1, R_2) = B_x(R_2) \backslash B_x(R_1)$, and $A = A(x, \frac{1}{3}R_2)$. Here $V_K^{n+4A}(B_x(r))$ denotes the volume of the ball in the model space M_K^{n+4A}, i.e., the simply connected space form of dimension $n + 4A$ with constant sectional curvature $-K$.

For model space $V_K^{n+4A}(B_x(r))$, if $K > 0$, this model space is the hyperbolic space; if $K = 0$, this model space is the Euclidean space. In any case, we have the following lower and upper estimates

$$\omega_{n+4A} \cdot r^{n+4A} \leq V_K^{n+4A}(B_x(r)) \leq \omega_{n+4A} \cdot r^{n+4A} e^{(n-1+4A)\sqrt{K}\,r}, \tag{2.2}$$

where ω_{n+4A} is the volume of the unit ball in $(n+4A)$-dimensional Euclidean space.

In [23], P. Wu and the author proved the local Neumann Poincaré inequality on complete smooth metric measure spaces.

Lemma 2.2. *Let $(M, g, e^{-f}dv)$ be an n-dimensional complete noncompact smooth metric measure space with*

$$\mathrm{Ric}_f \geq -(n-1)\,K$$

for some constant $K \geq 0$. Fix a point $o \in M$ and $R > 0$. For any $x \in B(o, R)$, we have

$$\int_{B_x(r)} |\varphi - \varphi_{B_x(r)}|^2 e^{-f} dv \leq c_1 e^{c_2 A + c_3(1+A)\sqrt{K}r} \cdot r^2 \int_{B_x(r)} |\nabla \varphi|^2 e^{-f} dv \tag{2.3}$$

for all $0 < r < R$ *and* $\varphi \in C^\infty(B_x(r))$, *where* $\varphi_{B_x(r)} := \int_{B_x(r)} \varphi d\mu / \int_{B_x(r)} d\mu$; $A = \sup_{y\in B_o(3R)} |f(y)|$, *and where the coefficient* c_i *are constants, depending only on the dimension* n.

We remark that the local f-Neumann Poincaré inequality and so called f-volume doubling property can imply a local Sobolev inequality and a mean value inequality for solutions to the f-heat equation on smooth metric measure spaces. Putting these inequalities together, we can furthermore prove Moser's Harnack inequality by using the De Giorgi-Nash-Moser theory. For detailed discussions, the interested readers can refer to [23].

In this paper, we are concerned with analytic and geometric results on $(M, g, e^{-f}dv)$ when $\mathrm{Ric}_f \geq -c\, r^{-2}(x)$ for some constant $c > 0$. Under this curvature assumption, Lemma 2.1 implies that

Proposition 2.3. *Let* $(M, g, e^{-f}dv)$ *be an* n*-dimensional complete noncompact smooth metric measure space. If*

$$\mathrm{Ric}_f \geq -c\, r^{-2}(x)$$

for some constant $c > 0$, *where* $r(x) := d(o, x)$ *is the distance function, then*

$$V_f(B_o(R)) \leq C\, R^{n+4A}$$

for all $R > 1$, *where* C *depends on* n, $A := \sup_{y\in B_o(R)} |f(y)|$ *and* $V_f(B_o(1))$.

Proof. Fixing a point $o \in M$ and $R > 1$, then we have $r(x) \geq R$ for any point $x \in M \setminus B_o(R)$. Since

$$\mathrm{Ric}_f \geq -c\, r^{-2}(x)$$

for some constant $c > 0$, then

$$\mathrm{Ric}_f \geq -c\, R^{-2}$$

for any $R > 1$. Using Lemma 2.1, letting $r_1 = R_1 = 0$, $r_2 = 1$, $R_2 = R$ and $x = o$, then

$$\frac{V_f(B_o(R))}{V_f(B_o(1))} \leq \frac{V_K^{n+4A}(B_o(R))}{V_K^{n+4A}(B_o(1))} \leq C(n, A)R^{n+4A},$$

where $A = \sup_{y\in B_o(R)} |f(y)|$. Note that we also used (2.2) in the above second inequality. Hence

$$V_f(B_o(R)) \leq C\, R^{n+4A}$$

for all $R > 1$, where C depends on n, A and $V_f(B_o(1))$. □

In [6], A. Grigor'yan and L. Saloff-Coste introduced a useful notion: *remote ball*. Recall that, for a fixed point $o \in M$, a ball $B(x, \rho)$ in M is called *remote ball* if

$$d(o, x) \geq 3\rho.$$

When $\mathrm{Ric}_f \geq -c\, r^{-2}(x)$, A. Grigor'yan and L. Saloff-Coste [6] showed that all remote balls in manifold M satisfy the volume doubling property, the Poincaré inequality and the parabolic Harnack inequality. In our case, for all remote balls $B(x,\rho)$, Lemma 2.2 can be restated as

Proposition 2.4. *Let $(M, g, e^{-f}dv)$ be an n-dimensional complete noncompact smooth metric measure space. Let $o \in M$ be a fixed point and $R > 0$. If*

$$\mathrm{Ric}_f \geq -c\, r^{-2}(x)$$

for some constant $c > 0$, where $r(x) := d(o,x)$ is the distance function, then for all remote balls $B(x,\rho)$, $x \in B(o,R)$, we have

$$\int_{B_x(\rho)} |\varphi - \varphi_{B_x(\rho)}|^2 e^{-f} dv \leq c_4 e^{c_5 \bar{A}} \cdot R^2 \int_{B_x(\rho)} |\nabla \varphi|^2 e^{-f} dv \tag{2.4}$$

for any $\varphi \in C^\infty(B_x(\rho))$, where $\varphi_{B_x(\rho)} := \int_{B_x(\rho)} \varphi e^{-f} dv / \int_{B_x(\rho)} e^{-f} dv$ and $\bar{A} = \sup_{y \in B_o(3R)} |f(y)|$. Here c_4 and c_5 are constants depending only on n.

3. Proof of Theorem 1.1

In this section, we first follow the argument of [4] to give a useful gradient estimate in the integral sense. The proof depends on the cut-off function technique and iterated procedure. Then we apply this estimate to prove Theorem 1.1.

Lemma 3.1. *Let $(M, g, e^{-f}dv)$ be an n-dimensional complete noncompact smooth metric measure space. For any a point $o \in M$ and $R > 0$, let u be a f-harmonic smooth function on $B(o, 2R)$, and*

$$\mathrm{diam}\, \partial B(o,r) \leq \epsilon r \quad \text{with} \quad \epsilon \in (0, 1/12)$$

for all $r \in [R, 2R]$. Assume that all remote balls $B(x,\rho)$, $x \in B(o, 2R)$, satisfies a local Poincaré inequality:

$$\int_{B_x(\rho)} |\varphi - \varphi_{B_x(\rho)}|^2 e^{-f} dv \leq c_4 e^{c_5 \bar{A}} \cdot R^2 \int_{B_x(\rho)} |\nabla \varphi|^2 e^{-f} dv \tag{3.1}$$

for any $\varphi \in C^\infty(B_x(\rho))$, where $\varphi_{B_x(\rho)} := \int_{B_x(\rho)} \varphi e^{-f} dv / \int_{B_x(\rho)} e^{-f} dv$ and $\bar{A} := \sup_{y \in B_o(6R)} |f(y)|$. Then we have

$$\int_{B_o(R)} |\nabla u|^2 e^{-f} dv \leq \delta^{\frac{1}{\epsilon}-1} \int_{B_o(2R)} |\nabla u|^2 e^{-f} dv, \tag{3.2}$$

where $\delta := \left(\frac{9 c_4 e^{c_5 \bar{A}}}{1 + 9 c_4 e^{c_5 \bar{A}}}\right)^{\frac{1}{6}}$, and $\bar{A} := \sup_{y \in B_o(6R)} |f(y)|$.

Proof. For any a point $o \in M$ and $R > 0$, suppose that $u \colon B(o, 2R) \to \mathbb{R}$ be a f-harmonic function and $c \in \mathbb{R}$ is a real number. Choosing

$$r \in [R + 3\epsilon R, 2R - 3\epsilon R],$$

where $\epsilon \in (0, 1/12)$, the diameter hypothesis in lemma implies that there exists some $x \in \partial B(o,r)$ such that

$$B(o, r+\epsilon R) \setminus B(o,r) \subset B(x, \epsilon R + \epsilon r).$$

Let $\eta(x)$ be a C^2 cut-off function with support in $B(o, r+\epsilon R)$. That is, $\eta(x) = 1$ if $x \in B(o,r)$; $\eta(x) = \frac{r+\epsilon R - d(o,x)}{\epsilon R}$ if $x \in B(o, r+\epsilon R) \setminus B(o,r)$; $\eta(x) = 0$ if $x \in M \setminus B(o, r+\epsilon R)$.

We easily observe that

$$|\nabla(\eta(u-c))|^2 = \eta^2 |\nabla(u-c)|^2 + 2\eta(u-c)\langle \nabla\eta, \nabla(u-c)\rangle + (u-c)^2|\nabla\eta|^2.$$

Integrating this identity with respect to the weighted measure $e^{-f}dv$, we have

$$\int_M |\nabla(\eta(u-c))|^2 = \int_M \eta^2|\nabla(u-c)|^2 + 2\eta(u-c)\langle\nabla\eta, \nabla(u-c)\rangle + (u-c)^2|\nabla\eta|^2.$$

Notice that

(3.3)

$$\begin{aligned}\int_M \eta^2|\nabla(u-c)|^2 + 2\eta(u-c)\langle\nabla\eta, \nabla(u-c)\rangle &= \int_M \langle\nabla((u-c)\eta^2), \nabla(u-c)\rangle \\ &= -\int_M (u-c)\eta^2\Delta_f(u-c) \\ &= 0,\end{aligned}$$

where we used integration by part with respect with to $e^{-f}dv$ and used the fact that u is f-harmonic. Therefore,

$$\int_M |\nabla(\eta(u-c))|^2 e^{-f}dv = \int_{B(o,r+\epsilon R)} (u-c)^2|\nabla\eta|^2 e^{-f}dv.$$

According to the definition of $\eta(x)$, by the above equality, we can get that

$$\begin{aligned}\int_{B(o,r)} |\nabla u|^2 e^{-f}dv &\le \int_{B(o,r+\epsilon R)} |\nabla(\eta(u-c))|^2 e^{-f}dv \\ &= \int_{B(o,r+\epsilon R)} (u-c)^2|d\eta|^2 e^{-f}dv \\ &\le \frac{1}{\epsilon^2 R^2}\int_{B(o,r+\epsilon R)\setminus B(o,r)} (u-c)^2 e^{-f}dv \\ &\le \frac{1}{\epsilon^2 R^2}\int_{B(x,\epsilon R+\epsilon r)} (u-c)^2 e^{-f}dv.\end{aligned}$$

For the right hand side of the above inequality, since $\epsilon \le 1/12$, this implies that the ball $B(x, \epsilon R + \epsilon r)$ is remote. Hence if choosing

$$c = u_{B(x,\epsilon(R+r))} = \frac{1}{\int_{B(x,\epsilon(R+r))} e^{-f}dv} \cdot \int_{B(x,\epsilon(R+r))} ue^{-f}dv,$$

then using the local f-Poincaré inequality (3.1) and the fact that $r+R \leq 3R$, we have that

$$\int_{B(o,r)} |\nabla u|^2 e^{-f} dv \leq 9c_4 e^{c_5 \bar{A}} \int_{B(x,3\epsilon R)} |\nabla u|^2 e^{-f} dv,$$

where $\bar{A} = \sup_{y \in B_o(6R)} |f(y)|$. Also noticing that

$$B(x, 3\epsilon R) \subset B(o, r + 3\epsilon R) \setminus B(o, r - 3\epsilon R),$$

hence we get

$$\begin{aligned}\int_{B(o,r-3\epsilon R)} |\nabla u|^2 e^{-f} dv &\leq \int_{B(o,r)} |\nabla u|^2 e^{-f} dv \\ &\leq 9c_4 e^{c_5 \bar{A}} \int_{B(o,r+3\epsilon R)\setminus B(o,r-3\epsilon R)} |\nabla u|^2 e^{-f} dv.\end{aligned}$$

That is, for all $r \in [R, R - 6\epsilon R]$, we have

$$\int_{B(o,r)} |\nabla u|^2 e^{-f} dv \leq \frac{9c_4 e^{c_5 \bar{A}}}{1 + 9c_4 e^{c_5 \bar{A}}} \int_{B(o,r+6\epsilon R)} |\nabla u|^2 e^{-f} dv.$$

Iterating this inequality, we finally get

$$\int_{B(o,R)} |\nabla u|^2 e^{-f} dv \leq \left(\frac{9c_4 e^{c_5 \bar{A}}}{1 + 9c_4 e^{c_5 \bar{A}}} \right)^N \int_{B(o,2R)} |\nabla u|^2 e^{-f} dv$$

provided that

$$N 6\epsilon R \leq R.$$

At last the desired result follows by choosing $\delta = \left(\frac{9c_4 e^{c_5 \bar{A}}}{1+9c_4 e^{c_5 \bar{A}}} \right)^{\frac{1}{6}}$. □

Now we are ready to prove Theorem 1.1 by using Lemma 3.1, similar to the proof of Theorem 2.1 in [4],

Proof of Theorem 1.1. Let $u \colon M \to \mathbb{R}$ be a f-harmonic function with polynomial growth of order ν, namely,

$$u(x) \leq C(1 + d(o, x))^{\nu}.$$

For $R >> 1$, we define

$$I_R := \int_{B(o,R)} |\nabla u|^2 e^{-f} dv$$

and

$$\epsilon(r) := \sup_{t \geq r} \frac{\rho(t)}{t},$$

where $\rho(t) := \sup_{x,y \in \partial B(o,t)} d(x, y)$.

To estimate I_R, we shall introduce a $C_0^2(M)$ cut off function $\xi(x)$, which satisfies

$$\xi(x) = \begin{cases} 1 & \text{in} \quad B(o,R), \\ \frac{2R-d(o,x)}{R} & \text{in} \quad B(o,2R)\setminus B(o,R), \\ 0 & \text{in} \quad M\setminus B(o,2R). \end{cases}$$

Hence we have

$$\begin{aligned} I_R &\le \int_{B(o,2R)} |\nabla(\xi u)|^2 e^{-f} dv \\ &= \int_{B(o,2R)} \Big[|u|^2|\nabla\xi|^2 + \xi^2|\nabla u|^2 + 2\xi u\langle\nabla\xi,\nabla u\rangle\Big] e^{-f} dv \\ &= \int_{B(o,2R)} |u|^2|\nabla\xi|^2 e^{-f} dv \\ &\le C R^{2\nu+n+4\bar{A}-2}, \end{aligned} \tag{3.4}$$

where C depends on n, $\bar{A} := \sup_{x\in B_o(3R)} |f(x)|$ and $V_f(B_o(1))$. Here in the third line of (3.4), we used the same proposition as (3.3); in the last line of (3.4), we used Proposition 2.3.

On the other hand, if we iterate the inequality (3.2) proved in Lemma 3.1, we can show that for all R such that $\epsilon(R) \le 1/12$:

$$I_R \le \delta^{-\ell+\sum_{j=0}^{\ell-1}\frac{1}{\epsilon(2^jR)}} I_{2^\ell R},$$

where $\delta = \left(\frac{9c_4e^{c_5\bar{A}}}{1+9c_4e^{c_5\bar{A}}}\right)^{\frac{1}{6}}$. Applying (3.4) to the right hand side of the above inequality yields

$$I_R \le C\cdot \exp\left\{\ell\left[\left(\frac{1}{\ell}\sum_{j=0}^{\ell-1}\frac{1}{\epsilon(2^jR)} - 1\right)\cdot\ln\delta + (2\nu+n+4\bar{A}-2)\cdot\ln 2\right]\right\}, \tag{3.5}$$

where in the above inequality, $\bar{A} := \sup_{x\in B_o(3R)} |f(x)|$, $\delta := \left(\frac{9c_4e^{c_5\bar{A}}}{1+9c_4e^{c_5\bar{A}}}\right)^{\frac{1}{6}}$ and $C := C\left(n,\nu,R,\bar{A},V_f(B_o(1))\right)$.

For all sufficiently large R, the Cesaro convergence theorem shows that

$$\lim_{\ell\to+\infty}\frac{1}{\ell}\sum_{j=0}^{\ell-1}\frac{1}{\epsilon\,(2^jR)} = +\infty.$$

Meanwhile, by our assumption:

$$\lim_{R\to\infty}\sup_{x\in B(o,R)}|f(x)| < +\infty,$$

we know that

$$0<\delta<1 \quad \text{and} \quad \ln\delta<0$$

for any R (even though $\ell \to +\infty$). Therefore, if $\ell \to +\infty$ in (3.5), then we conclude that

$$I_R = 0$$

for all sufficiently large R. Therefore u is constant. □

References

1. D. Bakry, M. Emery, *Diffusion hypercontractivitives*, in Séminaire de Probabilités XIX, 1983/1984, in: Lecture Notes in Math., vol. 1123, Springer-Verlag, Berlin, 1985, 177-206.
2. D. Bakry, Z.-M. Qian, *Some new results on eigenvectors via dimension, diameter and Ricci curvature*, Adv. Math. **155** (2000), 98–153.
3. K. Brighton, *A Liouville-type theorem for smooth metric measure spaces*, J. Geome. Anal. **23** (2013), 562–570.
4. G. Carron, *Harmonic functions on manifolds whose large sphere are small*, arXiv:math.DG/1503.05506.
5. N. Charalambous, Z. Lu, *The L^1 Liouville property on weighted manifolds*, Complex analysis and dynamical systems VI. Part 1, 65–79, Contemp. Math. **653**, Amer. Math. Soc., Providence, RI, 2015.
6. A. Grigor'yan, L. Saloff-Coste, *Stability results for Harnack inequalities*, Ann. Inst. Fourier(Grenoble) **55** (2005), 825–890.
7. R. Hamilton, *The formation of singularities in the Ricci flow*, Surveys in Differential Geom. **2** (1995), 7–136, International Press.
8. A. Kasue, *Harmonic functions of polynomial growth on complete Riemannian manifolds I*, Proc. Symp. Pure Math., Vol. **54**, part II, (1993), 281–290.
9. A. Kasue, *Harmonic functions of polynomial growth on complete Riemannian manifolds II*, J. Math. Soc. Japan, **47** (1995), 37–65.
10. P. Li, *Uniqueness of L^1 solutions for the Laplace equation and the heat equation on Riemannian manifolds*, J. Differential Geom. **20** (1984), 447–457.
11. P. Li, R. Schoen, *L^p and mean value properties of subharmonic functions on Riemannian manifolds*, Acta Math. **153** (1984), 279–301.
12. X.-D. Li, *Liouville theorems for symmetric diffusion operators on complete Riemannian manifolds*, J. Math. Pure. Appl. **84** (2005), 1295–1361.
13. X.-D. Li, *Hamilton's Harnack inequality and the W-entropy formula on complete Riemannian manifolds*, Stochastic Process. Appl. **126**, 4, (2016), 1264–1283.
14. J. Lott, *Some geometric properties of the Bakry-Émery-Ricci tensor*, Comment. Math. Helv. **78** (2003), 865–883.
15. O. Munteanu, J.-P. Wang, *Smooth metric measure spaces with nonnegative curvature*, Comm. Anal. Geom. **19** (2011), 451–486.
16. S. Pigola, M. Rimoldi, A. G. Setti, *Remarks on non-compact gradient Ricci solitons*, Math. Z. **268** (2011), 777–790.
17. G. Wei, W. Wylie, *Comparison geometry for the smooth metric measure spaces*, Proceedings of the 4th International Congress of Chinese Mathematicians, **2** (2007), 191–202.
18. G. Wei, W. Wylie, *Comparison geometry for the Bakry-Émery Ricci tensor*, J. Differential Geom. **83** (2009), 377–405.
19. J.-Y. Wu, *A note on the splitting theorem for the weighted measure*, Ann. Glob. Anal. Geom. **43** (2013), 287–298.
20. J.-Y. Wu, *L^p-Liouville theorems on complete smooth metric measure spaces*, Bull. Sci. Math. **138** (2014), 510–539.
21. J.-Y. Wu, *Elliptic gradient estimates for a weighted heat equation and applications*, Math. Z. **280** (2015), 451–468.

22. J.-Y. Wu, P. Wu, *Heat kernel on smooth metric measure spaces with nonnegative curvature*, Math. Ann. **362** (2015), no. 3-4, 717–742.
23. J.-Y. Wu, P. Wu, *Heat kernel on smooth metric measure spaces and applications*, Math. Ann. **365** (2016), 309–344.
24. J.-Y. Wu, P. Wu, *On L^p-Liouville property for smooth metric measure spaces*, arXiv:math.DG/1410.7305.
25. S.-T. Yau, *Harmonic functions on complete Riemannian manifolds*, Comm Pure Appl Math. **28** (1975), 201–228.
26. S.-T. Yau, *Some function-theoretic properties of complete Riemannian manifolds and their applications to geometry*, Indiana Univ. Math. J. **25** (1976), 659–670.

Department of Mathematics, Shanghai Maritime University, 1550 Haigang Avenue, Shanghai 201306, P. R. China

E-mail address: jywu81@yahoo.com